

REG-14635939

VAUZGE

NLM --- W1 IN214G (Gen): Film S02838: E-Journal free at producer site

US PATENT AND TRADEMARK OFFICE
SCIENTIFIC AND TECHNICAL INFO CTR
107 S. WEST STREET, PMB 803
ALEXANDRIA, VA 22314

ATTN:	SUBMITTED:	2009-03-05 15:34:31
PHONE: 571-272-2517	PRINTED:	2009-03-06 10:12:16
FAX: 571-272-0230	REQUEST NO.:	REG-14635939
E-MAIL: STIC-DOCS@uspto.gov	SENT VIA:	DOCLINE
	DOCLINE NO.:	26651776

REG	Copy	Journal
TITLE: INDIAN JOURNAL OF MEDICAL RESEARCH		
PUBLISHER/PLACE: Council Of Medical Research New Delhi		
VOLUME/ISSUE/PAGES: 1961 Sep;49():781-7 781-7		
DATE: 1961		
AUTHOR OF ARTICLE: DHALLA N;SASTRY M;MALHOTRA C		
TITLE OF ARTICLE: SOME IN VITRO EFFECTS OF HERPESTIS MONNIERA LINN O		
ISSN: 0971-5916		
OTHER NUMBERS/LETTERS: Unique ID.: 0374701 26651776 13335951		
SOURCE: PubMed		
MAX COST: \$4.00		
COPYRIGHT COMP.: Guidelines		
CALL NUMBER: W1 IN214G (Gen): Film S02838: E-Journal free at producer site		
REQUESTER INFO: 679798		
DELIVERY: E-mail Post to Web: STIC-DOCS@uspto.gov		
REPLY: Mail:		

KEEP THIS RECEIPT TO RECONCILE WITH BILLING STATEMENT

For problems or questions, contact NLM at http://wwwcf.nlm.nih.gov/ill/ill_web_form.cfm or phone 301-496-5511.
Include LIBID and request number.

NOTE:--THIS MATERIAL MAY BE PROTECTED BY COPYRIGHT LAW (TITLE 17, U.S. CODE)

SOME IN VITRO EFFECTS OF *HERPESTIS MONNIERA* LINN ON THE RESPIRATION OF RAT BRAIN*.

N. S. DHALLA, M. S. SASTRY, AND C. L. MALHOTRA.

(From the Department of Pharmacology & Therapeutics, Lady Hardinge Medical College, New Delhi.)

(Received for publication, April 7, 1961.)

THE plant *Herpestis monniera* (N.O. *Scrophulariaceae*) indigenous to India has been claimed to be a potent nervine tonic. It is being extensively used for various nervous disorders, such as insanity, epilepsy, hysteria, etc., in the Ayurvedic system of medicine (Chopra, 1933).

Bose and Bose (1931) claimed that the plant contained a very small amount of alkaloid 'brahmine' which was found to be highly toxic in frogs and guinea-pigs. Sastry, Dhalla and Malhotra (1959) isolated hersaponin and D-mannitol and also confirmed the presence of alkaloids. Malhotra and Das (1959) reported that the total extract of *Herpestis monniera* has cardiotonic, vasoconstrictor, sedative and neuro-muscular blocking actions. It was also found that hersaponin has cardiotonic action in frogs, sedative action in albino rats and guinea-pigs, and spasmodic action on certain smooth muscles. Recently, Malhotra, Das and Dhalla (1960) have evaluated the sedative action of hersaponin in mice and have established its hypnotic potentiating activity using barbiturates and ethanol. Hersaponin did not protect rats against electroshock and mice against metrazol seizures. It produced hypothermia in mice and reduced the amphetamine toxicity in aggregated mice but had no effect on metrazol toxicity. It has also been suggested that the central action of hersaponin simulate, those of the known tranquillizers—reserpine and chlorpromazine, having some properties of each. Prakash and Sirsi (1960) have established the psychotropic action of *Herpestis monniera* in facilitating the 'motor learning' in rats. Subsequently, Aithal and Sirsi (1961) have shown that the alcoholic extract of the drug exhibited tranquillizing effect on rats and dogs. The present paper deals with the effect of total extract, alkaloidal fraction and hersaponin isolated from *Herpestis monniera* on the respiratory activity *in vitro* of rat brain.

MATERIALS AND METHODS.

Albino rats of either sex weighing 120 g. to 150 g. were used. The animals were killed by decapitation and the brain was immediately removed, weighed, washed and transferred to a previously chilled glass tissue grinder containing Krebs Ringer phosphate buffer pH 7.3 (Umbreit, Burris and Stauffer, 1957). The brain homogenate was diluted with buffer so that each c.c. of homogenate contained 100 mg. of the fresh tissue.

Oxygen consumption was measured at 10 minute intervals by conventional Warburg manometric technique. The gas phase was air and the temperature of the

* Studies carried out under 'Drug Research Unit' with grant from Indian Council of Medical Research.

bath was 38°C. Each reaction vessel contained 1 c.c. of the rat brain homogenate and solvent control (0.3 c.c.) with or without drug. In those experiments where substrates were used, 0.3 c.c. of glucose (0.33 M), or 0.3 c.c. of sodium glutamate (2.0 M), or 0.3 c.c. of sodium succinate (0.33 M), all dissolved in phosphate buffer were added. In some experiments, 0.3 c.c. of 10^{-3} M lysergic acid diethyl amide (LSD 25) or 0.3 c.c. of 10^{-3} M 5-hydroxytryptamine (5-HT), in phosphate buffer was added. The total volume of the fluid in the flask was always made up to 3 c.c. with phosphate buffer. The central well of the flask contained 0.2 c.c. of 20 per cent KOH absorbed on filter paper.

The drug was tipped in from the side arm at zero time after fifteen minutes equilibration. All the measurements were performed in triplicate and the rate of oxygen uptake, Q_{O_2} for the first sixty minutes was calculated as μ l. of oxygen per hour per 100 mg. of the brain tissue (Shatzko and Sim, 1959; Starbuck and Helm, 1959). All the concentrations of drug refer to the final concentrations in reaction vessel.

The drugs used were *Herpestis monniera* extract and its constituents isolated as given below:—

Total extract: (Yield 5 per cent).—The hot alcoholic extract was concentrated to remove the solvent and freed from chlorophyll and salts.

Alkaloidal fraction: (Yield 0.05 per cent).—Above extract was digested with 1 per cent hot hydrochloric acid. The acid solution was freed from acidic material and the bases were extracted with chloroform after making alkaline with ammonium hydroxide.

Hersaponin: (Yield 0.02 per cent).—It was isolated by the method described earlier (Sastri, Dhalla and Malhotra, 1959).

The total extract was taken in phosphate buffer, while the alkaloidal fraction and hersaponin were dissolved in phosphate buffer containing 1 per cent propylene glycol which also served as control. The effect of solvent control was found to be insignificant and hence omitted from the text.

RESULTS.

The rates of oxygen consumption at different intervals by 100 mg. of the brain homogenate in phosphate buffer in the presence of various concentrations of total extract and alkaloidal fraction of *Herpestis monniera* are given in Graphs 1 and 2 respectively. Each reading is the average of 9 to 21 reaction vessels in several experiments for the same concentration of the drug and is within the variable range of ± 5 . The effect of various concentrations of the total extract and other constituents of *Herpestis monniera* on the respiration of rat brain are given in Table I. The values show the oxygen uptake as μ l. of oxygen, per hour per 100 mg. of the brain tissue and are the average of the number of experiments mentioned against each concentration.

In each experiment the same homogenate was used for the oxygen uptake with or without the drug. Variations in the respiratory rate have been calculated separately

The effect

A—
in each e
in each ex
for the sat
statistical
effect.


Tab
degrees o
that herse
on the res

ED₅₀
and Wile
(1 : 4,000
compare
times mor
tion (1 : 5

The
glucose (

GRAPH 1.

The effect of different concentrations of total extract of *Herpestis monniera* on oxygen uptake of rat brain homogenate in phosphate buffer.



A—solvent control; B, C, D, E, F and G—different concentrations of total extract.

in each experiment. Though the rate of oxygen uptake in the control flasks varied in each experiment, the percentage effect of the drugs showed very little variation for the same concentration. All the experimental results have been subjected to the statistical analysis and the standard error calculated is based on the percentage effect.


Table I shows that both the total extract and alkaloidal fraction had varying degrees of inhibitory action on the oxidative process. It is quite interesting to note that hersaponin, even in higher concentrations had no considerable depressing action on the respiration of brain *in vitro*.

ED₅₀ (with 10/20 confidence limits) calculated by the method of Litchfield and Wilcoxon (1949) was found to be 1:446 (1:1,020—1:229) and 1:2,291 (1:4,060—1:1,348) for the total extract and alkaloidal fraction, respectively. A comparison of the ED₅₀ shows that the effect of the alkaloidal fraction was five times more than that of the total extract. On the other hand, in the same concentration (1:500), hersaponin was about one third as effective as the total extract.

The oxygen uptake by 100 mg. of the rat brain homogenate in the presence of glucose (0.03 M), sodium glutamate (0.20 M), and sodium succinate (0.03 M) as

GRAPH 2.

The effect of different concentrations of alkaloidal fraction from *Herpestis monniera* on oxygen uptake of rat brain homogenate in phosphate buffer containing 1 per cent propylene glycol.



A—solvent control; B, C, D and E—different concentrations of alkaloidal fractions.

substrates with or without the alkaloidal fraction of *Herpestis monniera* is given in Table II. The percentage inhibition was least in sodium glutamate, moderate in sodium succinate, and marked in glucose. The approximate ratio of the inhibitory effect of alkaloidal fraction in their presence is 2 : 2 : 5, respectively.

The influence of LSD-25 (10^{-4} M) and 5-HT (10^{-3} M) on the respiration with or without the alkaloidal fraction and hersaponin has been recorded in Table III. It will be seen from the table that the inhibitory effect produced by the addition of LSD-25 to the Warburg vessel containing hersaponin was less than that of either LSD-25 or hersaponin. There was no change in the action of alkaloidal fraction in the presence of LSD-25. 5-HT, on the other hand, potentiated the action of hersaponin considerably, even though it had individually inhibitory action on the respiration of brain *in vitro*, while it did not exhibit any influence on the action of alkaloidal fraction.

DISCUSSION.

The present studies show that the total extract of *Herpestis monniera* had inhibitory action on the respiration of rat brain homogenate. The action was found to be mainly due to its alkaloidal content. The inhibitory effect of the alkaloidal

Effect

A.

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

1:

TABLE I.

Effect of total extract of *Herpestis monniera* and its constituents on the oxygen uptake of rat brain homogenate.

Concentration of drug.		QO ₂ without drug*.	QO ₂ with drug*.	Per cent inhibition (with S. E.)
A. Total extract :				
1: 10,000	(3)†	75.0	68.7	8.3±0.6
1: 5,000	(3)	75.0	62.1	14.7±0.6
1: 2,000	(4)	74.8	49.7	30.8±2.0
1: 500	(4)	78.3	37.5	52.4±1.0
1: 100	(10)	78.8	23.5	67.8±1.0
1: 20	(5)	78.8	9.2	87.1±1.2
B. Alkaloidal fraction :				
1: 10,000	(3)	77.0	66.0	13.6±2.0
1: 5,000	(4)	67.6	44.6	32.3±0.6
1: 3,000	(6)	68.9	38.3	43.2±1.4
1: 2,000	(5)	73.1	32.3	55.1±1.8
C. Hersaponin :				
1: 5,000	(4)	74.9	72.1	3.8±0.5
1: 1,000	(5)	78.7	70.5	7.9±1.8
1: 500	(3)	71.2	58.7	16.1±1.2

*Calculated as $\mu\text{L/hr./100 mg. of brain tissue.}$

†Figures within brackets show the number of experiments carried out in each study.

TABLE II.

Effect of alkaloidal fraction (5×10^{-4}) with or without various substrates-glucose, sodium glutamate and sodium succinate on the respiration of rat brain.

Substrate.		QO ₂ without drug*.	QO ₂ with drug*.	Per cent inhibition. (with S.E.).
Control	(5)†	74.2	33.7	54.5±1.5
Glucose (0.02M)	(5)	60.6	40.2	50.1±1.7
Sodium glutamate (0.02M)	(4)	53.4	67.9	18.6±1.1
Sodium succinate (0.02M)	(4)	110.2	80.1	27.4±2.3

*Calculated as $\mu\text{L/hr./100 mg. of brain tissue.}$

†Figures within brackets show the number of experiments carried out in each study.

fraction on the oxidative process could not be significantly influenced by 5-HT or LSD-25. However, there was slight potentiation of action in each case. From our study of the influence of substrates on the action of alkaloidal fraction, it is quite apparent that both sodium glutamate and sodium succinate partially block its inhibitory effect. It is, therefore, suggested that the alkaloidal fraction is suppressing the enzyme system and the action might be localized somewhere in the citric acid cycle. Further work is being carried out to elucidate the mechanism of action of alkaloidal fraction *in vivo*.

Malhotra, Das and Dhalla (1960) have reported that the hypnotic potentiating action of hersaponin was inhibited on pretreatment with LSD-25 and also have

TABLE III.

Effect of lysergic acid diethyl amide and 5-hydroxytryptamine with or without alkaloidal fraction (5×10^{-4}) and hersaponin (1×10^{-3}) on the respiration of rat brain.

Drug.	Q_{O_2} in μ l./hr./100 mg. of brain tissue*.	Per cent inhibition. (with S.E.).	Probability†.
Control	74.3		..
LSD-25 (10^{-4} M)	65.8	11.3 ± 0.4	..
5-HT (10^{-4} M)	58.3	21.9 ± 0.8	..
Alkaloidal fraction	34.8	53.7 ± 1.2	..
Alkaloidal fraction plus LSD-25 (10^{-4} M)	31.5	57.8 ± 1.4	$P > 0.05$
Alkaloidal fraction plus 5-HT (10^{-4} M)	32.0	56.8 ± 1.2	$P > 0.05$
Hersaponin	67.1	10.0 ± 0.2	..
Hersaponin plus LSD-25 (10^{-4} M)	70.2	5.3 ± 0.1	$P < 0.05$
Hersaponin plus 5-HT (10^{-4} M)	48.9	34.7 ± 1.0	$P < 0.001$

*Four experiments were carried out in each study. †Probability (P) of no significance calculated by 't' test.

suggested that release of 5-HT might be one of the ways in which hersaponin acts. From our observations, it is evident that LSD-25 reduced the respiratory inhibition caused by hersaponin and is in agreement with *in vivo* results. Potentiation caused by 5-HT *in vitro* on the inhibiting action of hersaponin lends further support to the suggested mechanism *in vivo*. Since hersaponin exerted very slight respiratory inhibition *in vitro*, it is suggested that the marked sedative and hypnotic potentiating effects of hersaponin might be due to the formation of some metabolic product in the body.

Substances, such as 2, 4-Dinitrophenol which depress the formation of high energy phosphate bond without markedly affecting oxygen consumption are known as uncoupling agents. Reserpine and chlorpromazine are also found to inhibit oxidative phosphorylation (Kirpekar and Lewis, 1960). As this very new drug 'hersaponin' has been shown to possess great potentialities with special reference to central nervous system, it would therefore, be interesting to study its influence on uncoupling oxidative phosphorylation.

SUMMARY.

1. The effect of the total extract of *Herpestis monniera* and its constituents, namely alkaloidal fraction and hersaponin on the oxygen uptake of the rat brain homogenate has been studied by Warburg technique.

2. Total extract and alkaloidal fraction inhibited the respiration of brain tissue to a varying degree. ED_{50} were found to be 1 : 446 and 1 : 2,201, respectively. Hersaponin had a mild inhibitory action.

3. In the presence of glucose, sodium glutamate and sodium succinate as substrates, the alkaloidal fraction in concentration of 5×10^{-4} exhibited 50.1, 18.6 and 27.4 per cent inhibition, respectively.

4. LSD fraction, when potentiated by

Authors' investigation. the supply of

ARTHA, H.

BOSE, K.C.

CHOPRA, R.

KIRPEKAR, S.

LITCHFIELD (1940)

MALHOTRA,

MALHOTRA, N.S. (1960)

PRASAD, J.

SANTRY, M. C.L. (1960)

SHATZKO, P.

STARBUCK,

UMBERT, J. FER, J.

with or without
respiration of

1.	Probabilität.
----	---------------

1
 2
 3
 4
 5
 6
 7
 8
 9
 10
 11
 12
 13
 14
 15
 16
 17
 18
 19
 20
 21
 22
 23
 24
 25
 26
 27
 28
 29
 30
 31
 32
 33
 34
 35
 36
 37
 38
 39
 40
 41
 42
 43
 44
 45
 46
 47
 48
 49
 50
 51
 52
 53
 54
 55
 56
 57
 58
 59
 60
 61
 62
 63
 64
 65
 66
 67
 68
 69
 70
 71
 72
 73
 74
 75
 76
 77
 78
 79
 80
 81
 82
 83
 84
 85
 86
 87
 88
 89
 90
 91
 92
 93
 94
 95
 96
 97
 98
 99
 100
 101
 102
 103
 104
 105
 106
 107
 108
 109
 110
 111
 112
 113
 114
 115
 116
 117
 118
 119
 120
 121
 122
 123
 124
 125
 126
 127
 128
 129
 130
 131
 132
 133
 134
 135
 136
 137
 138
 139
 140
 141
 142
 143
 144
 145
 146
 147
 148
 149
 150
 151
 152
 153
 154
 155
 156
 157
 158
 159
 160
 161
 162
 163
 164
 165
 166
 167
 168
 169
 170
 171
 172
 173
 174
 175
 176
 177
 178
 179
 180
 181
 182
 183
 184
 185
 186
 187
 188
 189
 190
 191
 192
 193
 194
 195
 196
 197
 198
 199
 200
 201
 202
 203
 204
 205
 206
 207
 208
 209
 210
 211
 212
 213
 214
 215
 216
 217
 218
 219
 220
 221
 222
 223
 224
 225
 226
 227
 228
 229
 230
 231
 232
 233
 234
 235
 236
 237
 238
 239
 240
 241
 242
 243
 244
 245
 246
 247
 248
 249
 250
 251
 252
 253
 254
 255
 256
 257
 258
 259
 260
 261
 262
 263
 264
 265
 266
 267
 268
 269
 270
 271
 272
 273
 274
 275
 276
 277
 278
 279
 280
 281
 282
 283
 284
 285
 286
 287
 288
 289
 290
 291
 292
 293
 294
 295
 296
 297
 298
 299
 300
 301
 302
 303
 304
 305
 306
 307
 308
 309
 310
 311
 312
 313
 314
 315
 316
 317
 318
 319
 320
 321
 322
 323
 324
 325
 326
 327
 328
 329
 330
 331
 332
 333
 334
 335
 336
 337
 338
 339
 340
 341
 342
 343
 344
 345
 346
 347
 348
 349
 350
 351
 352
 353
 354
 355
 356
 357
 358
 359
 360
 361
 362
 363
 364
 365
 366
 367
 368
 369
 370
 371
 372
 373
 374
 375
 376
 377
 378
 379
 380
 381
 382
 383
 384
 385
 386
 387
 388
 389
 390
 391
 392
 393
 394
 395
 396
 397
 398
 399
 400
 401
 402
 403
 404
 405
 406
 407
 408
 409
 410
 411
 412
 413
 414
 415
 416
 417
 418
 419
 420
 421
 422
 423
 424
 425
 426
 427
 428
 429
 430
 431
 432
 433
 434
 435
 436
 437
 438
 439
 440
 441
 442
 443
 444
 445
 446
 447
 448
 449
 450
 451
 452
 453
 454
 455
 456
 457
 458
 459
 460
 461
 462
 463
 464
 465
 466
 467
 468
 469
 470
 471
 472
 473
 474
 475
 476
 477
 478
 479
 480
 481
 482
 483
 484
 485
 486
 487
 488
 489
 490
 491
 492
 493
 494
 495
 496
 497
 498
 499
 500
 501
 502
 503
 504
 505
 506
 507
 508
 509
 510
 511
 512
 513
 514
 515
 516
 517
 518
 519
 520
 521
 522
 523
 524
 525

- significance calculated

formation of high
hydrogen consumption
are also found to
As this very new
ilities with special
resting to study its

d its constituents,
ake of the rat brain

expiration of brain
2,201, respectively.

sodium succinate as
sites 50.1, 18.6 and